



National  
Library  
of Medicine

|        |            |   |  |  |                           |                      |                       |
|--------|------------|---|--|--|---------------------------|----------------------|-----------------------|
| PubMed | Nucleotide | Protein   | Genome   | Structure  | PopSet                    | Taxonomy             | OMIM                  |
| Search | PubMed     | <input style="width: 100px; border: 1px solid black;" type="text"/> for                             | <input style="border: 1px solid black; padding: 2px 5px;" type="button" value="Go"/> <input style="border: 1px solid black; padding: 2px 5px;" type="button" value="Clear"/> |  |                           |                      |                       |
|        |            | <a href="#">Limits</a>  | <a href="#">Preview/Index</a>  | <a href="#">History</a>  | <a href="#">Clipboard</a> |                      |                       |
|        |            | <a href="#">Display</a>   | <a href="#">Abstract</a>   | <input style="border: 1px solid black; width: 15px; height: 15px;" type="button" value="▼"/> | <a href="#">Save</a>      | <a href="#">Text</a> | <a href="#">Order</a> |
|        |            | <input style="border: 1px solid black; padding: 2px 10px;" type="button" value="Add to Clipboard"/> |  |  |                           |                      |                       |

Entrez PubMed

PubMed Services

Related Resources

1: *Int J Dermatol* 1998 Aug;37(8):595-8 [Related Articles](#), [Books](#), [LinkOut](#)

## A new model of epidermal culture for the surgical treatment of vitiligo.

**Andreassi L, Pianigiani E, Andreassi A, Taddeucci P, Biagioli M.**

Department of Dermatology, University of Siena, Italy.

**BACKGROUND:** Vitiligo can be successfully treated with grafts of autologous cultured epidermal cells. **OBJECTIVE:** To evaluate the efficacy of autologous grafting of epidermal cells, cultured by an original method, in the treatment of localized vitiligo refractory to other therapies. **METHODS:** Autologous normally pigmented skin was used to culture keratinocytes and melanocytes on a supporting layer of biomaterial (Laserskin), which was grafted directly onto achromatic skin after de-epithelialization with liquid carbon dioxide. The percentage area of repigmentation was calculated by image analysis. **RESULTS:** Initial repigmentation of the treated areas was observed 1 month after treatment. Repigmentation continued to increase for 3 months after grafting. Follow-up at 3, 6, 12, and 18 months showed almost complete repigmentation in six out of 11 cases. In four other patients, 40-71% of the grafted achromatic area was repigmented. In one patient, repigmentation was impeded by sepsis. **CONCLUSIONS:** The method was found to be effective in the treatment of localized vitiligo refractory to other treatments. The therapeutic procedure was simple, reproducible, and easy to use.

PMID: 9732005 [PubMed - indexed for MEDLINE]

|         |          |  |      |      |       |   |
|---------|----------|--|------|------|-------|---|
| Display | Abstract | <input style="border: 1px solid black; width: 15px; height: 15px;" type="button" value="▼"/> | Save | Text | Order | <input style="border: 1px solid black; padding: 2px 10px;" type="button" value="Add to Clipboard"/> |
|---------|----------|--|------|------|-------|---|

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)  
[Freedom of Information Act](#) | [Disclaimer](#)

[Freedom of Information Act](#) | [Disclaimer](#)